

MULTI-ANTIGEN IMAGING REVEALS INFLAMMATORY DC, ADAM17 AND NEPRILYSIN AS EFFECTORS IN KELOID FORMATION

Jürgen Bauerschmitz ^{3,4,5} Mathias Rath ¹, Alain Pitiot ², Michael Kirr ^{3,4,5}, Waltraud Fröhlich ^{3,4,5}, Bianca Plosnita ⁶, Stefan Schliep ^{3,4,5}, Andreas S Baur ^{3,4,5}, Christian Ostalecki ³

Jürgen Bauerschmitz ^{3,4,5} Mathias Rath ¹,
Alain Pitiot ², Michael Kirr ^{3,4,5}, Waltraud
Fröhlich ^{3,4,5}, Bianca Plosnita ⁶, Stefan
Schliep ^{3,4,5}, Andreas S Baur ^{3,4,5}, Christian
Ostalecki ³

¹Department of Urology, University Hospital Heidelberg, 69120 Heidelberg, Germany.

²Laboratory of Image & Data Analysis, Ilixa Ltd., London W1U 6NQ, UK.

³Department of Dermatology, Universitätsklinikum Erlangen, Friedrich-Alexander-University Erlangen-Nürnberg, 91054 Erlangen, Germany.

⁴Deutsches Zentrum Immuntherapie (DZI), 91054 Erlangen, Germany.

⁵Comprehensive Cancer Center (CCC) Erlangen, 91054 Erlangen, Germany.

⁶TissueGnostics GmbH, 1020 Vienna, Austria.

⁷Department of Psychiatry and Psychotherapy, Universitätsklinikum Erlangen, Friedrich-Alexander-University Erlangen-Nürnberg, 91054 Erlangen, Germany.

Running Title

Multi-Antigen Imaging Reveals Inflammatory DC, ADAM17 and Neprilysin as Effectors in Keloid Formation

Word Count - 213 words

BACKGROUND

Keloid disorder is an aberrant scarring process of the skin, characterized by excessive extracellular matrix synthesis and deposition. The pathogenesis of this prevalent cutaneous disorder is not fully understood; however, a persistent inflammatory process is observed. To obtain more insight into this process, we analyzed lesional, perilesional and healthy tissue using multi-antigen-analysis (MAA) in conjunction with a data mining approach.

METHODS

To obtain more insight into this process, we analyzed lesional, perilesional and healthy tissue using multi-antigen-analysis (MAA) in conjunction with a data mining approach.

RESULTS

Here, we demonstrate that monocyte-derived inflammatory dendritic cells (CD1a+, CD11c+, CD14+) and activated CD4+ T lymphocytes (CD45 RO+) dominated the immune infiltration in keloids while associating with fibroblasts. In perilesional tissue, precursor immune cells were dominant in the perivascular area, suggesting that they were attracted by an immune process, potentially in the lesional area. Supporting this hypothesis, only in keloid lesions, high levels of ADAM10/17 and Neprilysin (CD10) were observed in both fibroblasts and leukocytes. The spatial proximity of these two cell types, which could be confirmed by image analysis only in lesional tissue, could be a potential factor leading to the activation of fibroblasts.

CONCLUSION

Our findings provide new insight into the pathogenesis of keloid formation and reveal metalloproteinases as a target for therapeutical intervention.